

THREE THEOREMS ON TWIN PRIMES

VLADIMIR SHEVELEV

ABSTRACT. For earlier considered sequence: $c(1) = 2$ and for $n \geq 2$,

$$c(n) = c(n-1) + \begin{cases} \gcd(n, c(n-1)), & \text{if } n \text{ is even} \\ \gcd(n-2, c(n-1)), & \text{if } n \text{ is odd} \end{cases}$$

we prove theorems of its connection with twin primes. We also give a sufficient condition for the infinity of twin primes and pose several new conjectures; among them we propose a very simple conjectural algorithm of constructing a pair $(p, p+2)$ of twin primes over arbitrary given integer $m \geq 4$ such that $p+2 \geq m$.

1. INTRODUCTION

In [2] we posed the following conjecture

Conjecture 1. *Let $c(1) = 2$ and for $n \geq 2$,*

$$c(n) = c(n-1) + \begin{cases} \gcd(n, c(n-1)), & \text{if } n \text{ is even} \\ \gcd(n-2, c(n-1)), & \text{if } n \text{ is odd} \end{cases}.$$

Then every record (more than 3) of the values of difference $c(n) - c(n-1)$ is greater of twin primes.

The first such records are (cf. sequence A166945 in [4])

$$(1.1) \quad 7, 13, 43, 139, 313, 661, 1321, 2659, 5459, 10891, 22039, \dots$$

Our observations of the behavior of sequence $\{c(n)\}$ are the following:

- 1) In some sequence of arguments $\{m_i\}$ we have $\frac{c(m_i)}{m_i} = 2$. These values of arguments we call *the fundamental points*. The least fundamental point is $m_1 = 2$.
- 2) For every two adjacent fundamental points $m_j < m_{j+1}$, we have $m_{j+1} \geq 2m_j$.
- 3) For $i \geq 2$, the numbers $m_i \mp 1$ are twin primes (and, consequently, $m_i \equiv 0 \pmod{6}$).
- 4) In points $m_i + 3$ we have $c(m_i + 3) - c(m_i + 2) = m_i + 1$. These increments we call *the main increments* of sequence $\{c(n)\}$, while other nontrivial (i.e. more than 1) increments we call *the minor increments*.

- 5) For $i \geq 2$, denote h_i the number of minor increments between adjacent fundamental points m_i and m_{i+1} and T_i the sum of these increments. Then $T_i \equiv h_i \pmod{6}$.
- 6) For $i \geq 3$, the minor increments between adjacent fundamental points m_i and m_{i+1} could occur only before $m_{i+1} - \sqrt{m_{i+1} - 1} - 4$.

The aim of this paper is to show that the validity of all these observations follow only from 6).

Theorem 1. *If observation 6) is true then observation 1)-5) are true as well.*

Corollary 1. *If 1) observation 6) is true and 2) the sequence $\{c(n)\}$ contains infinitely many fundamental points, then there exist infinitely many twin primes.*

Besides, in connection with Conjecture 1 we think that

Conjecture 2. *For $n \geq 1$, the main and only main increments are the record differences $c(n) - c(n-1)$.*

2. PROOF OF THEOREM 1

Note that $c(12) = 24$ and the numbers 11, 13 are primes. We use induction. Suppose $n_1 \geq 15$ is a number of the form $6l+3$ (for $n_1 < 15$ the all observations are verified directly). Let $n_1 - 3$ is a fundamental point:

$$c(n_1 - 3) = 2n_1 - 6$$

and for $n := n_1 - 3$, $n \mp 1$ are twin primes. Since $n_1 - 2$ is odd, then we have

$$c(n_1 - 2) = 2n_1 - 5.$$

Further, since $\gcd(n_1 - 1, 2n_1 - 5) = \gcd(n_1 - 1, 3) = 1$, then

$$c(n_1 - 1) = 2n_1 - 4,$$

Since

$$\gcd(n_1 - 2, 2n_1 - 4) = n_1 - 2$$

then we have a main increment such that

$$(2.1) \quad c(n_1) = 3n_1 - 6.$$

Here we distinguish two cases:

A) Up to the following fundamental point there are only trivial increments. The inductive step in this case we formulate as the following.

Theorem 2. *If $6 \leq m_j < m_{j+1}$ are adjacent fundamental points with only nontrivial increment between them which is a main increment then*

- i) $m_{j+1} = 2m_j$;
 ii) If $m_j \mp 1$ are twin primes, then $m_{j+1} \mp 1$ are twin primes as well.

Thus, if observation 2) is true, then to every pair of the adjacent fundamental points with the only main increment between them corresponds a quadruple of primes of the form

$$p, \quad p + 2, \quad 2p + 1, \quad 2p + 3.$$

Example 1. Consider the adjacent fundamental points $n_1 = 660$ and $n_2 = 1320$. Since $n_2 = 2n_1$, then between them there is no any miner increment. We have

$$c(660) = 1320, \quad c(661) = 1321, \quad c(662) = 1322, \quad c(663) = 1983$$

and

$$c(663) - c(662) = 661.$$

$$c(1320) = 2640, \quad c(1321) = 2641, \quad c(1322) = 2642, \quad c(1323) = 3963$$

and

$$c(1323) - c(1322) = 1321.$$

Here we have two pairs of twins:

$$p = 659, \quad p + 2 = 661, \quad 2p + 1 = 1319, \quad 2p + 3 = 1321.$$

Inductive step in case A)

Continuing (2.1), we have

$$c(n_1 + 1) = 3n_1 - 5,$$

$$c(n_1 + 2) = 3n_1 - 4,$$

...

$$c(2n_1 - 6) = 4n_1 - 12,$$

(It is the second fundamental point in the inductive step)

$$c(2n_1 - 5) = 4n_1 - 11,$$

$$c(2n_1 - 4) = 4n_1 - 10,$$

Since

$$\gcd(2n_1 - 5, 4n_1 - 10) = 2n_1 - 5$$

then, denoting $n_2 = 2n_1 - 3$, we have

$$(2.2) \quad c(2n_2) = 3n_2 - 6.$$

Note that, since $n_1 = 6l + 3$, then $n_2 = 6l_1 + 3$, where $l_1 = 2l$.

Furthermore, from the run of formulas (2.2) we find for $5 \leq j \leq \frac{n_1-2}{2}$

$$c(2n_1 - 2j - 1) = 4n_1 - 2j - 7,$$

$$c(2n_1 - 2j) = 4n_1 - 2j - 6.$$

This means that

$$\gcd(2n_1 - 2j, 4n_1 - 2j - 7) = 1, \text{ i.e. } \gcd(2j - 7, 2n_1 - 7) = 1.$$

Note that, for the considered values of n_1 we have $2\frac{n_1-2}{2} - 7 \geq \sqrt{2n_1 - 7}$, then $2n_1 - 7 = n_2 - 4$ is prime.

On the other hand,

$$c(2n_1 - 2j) = 4n_1 - 2j - 6,$$

$$c(2n_1 - 2j + 1) = 4n_1 - 2j - 5.$$

Thus, for $5 \leq j \leq \frac{n_1-1}{2}$,

$$\gcd(2n_1 - 2j - 1, 4n_1 - 2j - 6) = 1, \text{ i.e. } \gcd(j - 2, 2n_1 - 5) = 1.$$

Here, for the considered values of n_1 we also have $\frac{n_1-5}{2} \geq \sqrt{2n_1 - 5}$, then $2n_1 - 5 = n_2 - 2$ is prime as well. This completes the inductive step in case **A**). If, in addition, to note that $n_1 - 3$ and $n_2 - 3$ are the two adjacent fundamental points, then we get a proof of Theorem 2. ■

B) Up to the following fundamental point we have some minor increments.

The inductive step we formulate as following.

Theorem 3. *Let observation 6) be true. If $6 \leq m_i < m_{i+1}$ are adjacent fundamental points with a finite number of minor increments between them, then*

$$i) \ m_{i+1} \geq 2m_i;$$

$$ii) \ \text{If } m_i \mp 1 \text{ are twin primes, then } m_{i+1} \mp 1 \text{ are twin primes as well.}$$

Thus the observation 2) will be proved in frameworks of the induction.

Inductive step in case **B**)

Let in the points $n_1 + l_j$ $j = 1, \dots, h$, before the second fundamental point we have the minor increments t_j , $j = 1, \dots, h$. We have (starting with the first fundamental point $n_1 - 3$)

$$c(n_1 - 3) = 2n_1 - 6$$

$$c(n_1 - 2) = 2n_1 - 5,$$

$$c(n_1 - 1) = 2n_1 - 4,$$

$$c(n_1) = 3n_1 - 6,$$

$$\begin{aligned}
& c(n_1 + 1) = 3n_1 - 5, \\
& \dots \\
& c(n_1 + l_1 - 1) = 3n_1 + l_1 - 7. \\
(2.3) \quad & c(n_1 + l_1) = 3n_1 + l_1 + t_1 - 7, \\
& c(n_1 + l_1 + 1) = 3n_1 + l_1 + t_1 - 6, \\
& \dots
\end{aligned}$$

$$\begin{aligned}
& c(n_1 + l_2 - 1) = 3n_1 + l_2 + t_1 - 8, \\
(2.4) \quad & c(n_1 + l_2) = 3n_1 + l_2 + t_1 + t_2 - 8, \\
& \dots \\
& c(n_1 + l_h - 1) = 3n_1 + l_h + t_1 + \dots + t_{h-1} - h - 7, \\
(2.5) \quad & c(n_1 + l_h) = 3n_1 + l_h + t_1 + \dots + t_h - h - 6, \\
& c(n_1 + l_h + 1) = 3n_1 + l_h + t_1 + \dots + t_h - h - 5, \\
& \dots
\end{aligned}$$

$$(2.6) \quad c(2n_1 + T_h - h - 6) = 4n_1 + 2T_h - 2h - 12,$$

where

$$(2.7) \quad T_h = t_1 + \dots + t_h.$$

(thus $2n_1 + T_h - h - 6$ is the second fundamental point in the inductive step)

$$c(2n_1 + T_h - h - 5) = 4n_1 + 2T_h - 2h - 11,$$

$$c(2n_1 + T_h - h - 4) = 4n_1 + 2T_h - 2h - 10.$$

Here we need a lemma.

Lemma 1. $T_h - h$ is even.

Proof. We use the induction over $h \geq 1$. If l_1 is even, then, by (2.3),

$$\gcd(n_1 + l_1 - 2, 3n_1 + l_1 - 7) = t_1$$

and t_1 divides $2n_1 - 5$. Analogously, if l_1 is odd, then t_1 divides $2n_1 - 7$.

Thus $T_1 - 1 = t_1 - 1$ is even. Suppose that $T_{h-1} - (h - 1)$ is even. Then, by (2.5) in the case of l_h is even, we have

$$\gcd(n_1 + l_h - 2, 3n_1 + l_h + T_{h-1} - (h - 1) - 7) = t_h$$

and t_h divides, by the inductive supposition, an odd number $2n_1 + T_{h-1} - (h - 1) - 5$. Analogously, if l_h is odd, then t_h divides odd number $2n_1 + T_{h-1} - (h - 1) - 7$. Thus t_h is odd and we conclude that $T_h - h = T_{h-1} - (h - 1) + t_h - 1$ is even. ■

Therefore, we have

$$\gcd(2n_1 + T_h - h - 5, 4n_1 + 2T_h - 2h - 10) = 2n_1 + T_h - h - 5,$$

and in the point $n_2 := 2n_1 + T_h - h - 3$ we have the second main increment (in framework of the inductive step). Thus

$$(2.8) \quad c(2n_1 + T_h - h - 3) = 6n_1 + 3T_h - 3h - 15.$$

Note that, for $n \geq 2$, we have $c(n) \equiv n \pmod{2}$. Therefore, $T_h \geq 3h$ and for the second fundamental point $n_2 - 3 = 2n_1 + T_h - h - 6$ we find

$$n_2 - 3 \geq 2(n_1 - 3) + 2h.$$

By the induction (with Theorem 2), *this proves observation 2*).

Now, in order to finish the induction, we prove the primality of numbers $n_2 - 4 = 2n_1 + T_h - h - 7$ and $n_2 - 2 = 2n_1 + T_h - h - 5$.

From the run of formulas (2.5)-(2.6) for $5 \leq j \leq \frac{n_1 + T_h - h - l_h - 3}{2}$ (unfortunately, we cannot cross the upper boundary of the last miner increment) we find

$$c((2n_1 + T_h - h - 4) - (2j - 1)) = 4n_1 + 2T_h - 2h - 2j - 9,$$

$$c(2n_1 + T_h - h - 2j - 2) = 4n_1 + 2T_h - 2h - 2j - 8.$$

This means that

$$\gcd(2n_1 + T_h - h - 2j - 2, 4n_1 + 2T_h - 2h - 2j - 9) = 1,$$

i.e.

$$\gcd(2j - 7, 2n_1 + T_h - h - 7) = 1.$$

For the most possible $j = \frac{n_1 + T_h - h - l_h - 3}{2}$ we should have

$$2j - 7 = n_1 + T_h - h - l_h - 10 \geq \sqrt{2n_1 + T_h - h - 7},$$

or, since $n_2 = 2n_1 + T_h - h - 3$, then we should have $n_2 - n_1 - l_h - 7 \geq \sqrt{2n_1 + T_h - h - 7}$, i.e.

$$n_1 + l_h \leq n_2 - \sqrt{n_2 - 4} - 7.$$

This condition is equivalent to the observation 6) which is written in

terms of the fundamental points $m_i = n_i - 3$. Thus from observation 6) we indeed obtain the primality of $n_2 - 4 = 2n_1 + T_h - h - 7$.

What is left-to prove the primality of $n_2 - 2 = 2n_1 + T_h - h - 5$. We do it in the next section *without supposition of the validity of observation 6)*.

3. COMPLETION OF PROOF OF THEOREM 1: PROOF OF THE PRIMALITY OF $2n_1 + T_h - h - 5$ INDEPENDENTLY ON OBSERVATION 6)

It is interesting that, using the Rowland's method [1], we are able to get the primality of $2n_1 + T_h - h - 5$ without unproved observation 6). This gives an additional hope to convert the observations 1)-6) into the absolute statements.

Denote

$$(3.1) \quad n_1^* := n_1 + l_h.$$

By (2.5),

$$(3.2) \quad c(n_1^*) = 3n_1 + l_h + T_h - h - 6 = 3n_1^* - 2l_h + T_h - h - 6 = 3n_1^* + u,$$

where

$$(3.3) \quad u = T_h - h - 2l_h - 6 \equiv 0 \pmod{2},$$

and

$$(3.4) \quad c(n_1^* + i - 1) = 3n_1^* + i + u - 1, \quad i \leq k,$$

where k is the smallest positive integer such that the point $3n_1^* + k + u - 1$ is point of a nontrivial increment. Put $h(n) = c(n) - c(n - 1)$, such that

$$(3.5) \quad h(n) = \begin{cases} \gcd(n, c(n - 1)), & \text{if } n \text{ is even,} \\ \gcd(n - 2, c(n - 1)), & \text{if } n \text{ is odd,} \end{cases}$$

then

$$h(n_1^* + i) = \begin{cases} \gcd(n_1^* + i, c(n_1^* + i - 1)), & \text{if } n_1^* + i \text{ is even,} \\ \gcd(n_1^* + i - 2, c(n_1^* + i - 1)), & \text{if } n_1^* + i \text{ is odd.} \end{cases}$$

Put

$$(3.6) \quad \delta = \delta(m) = \begin{cases} 0, & \text{if } m \text{ is even,} \\ 2, & \text{if } m \text{ is odd.} \end{cases}$$

Thus, $h(n_1^* + i)$ divides both $n_1^* + i - \delta(n_1^* + i)$ and $3n_1^* + i + u - 1$ and also divides both their difference

$$(3.7) \quad 2n_1^* + u - 1 + \delta(n_1^* + i)$$

and

$$(3.8) \quad 3(n_1^* + i - \delta(n_1^* + i)) - (3n_1^* + i + u - 1) = 2i - u + 1 - 3\delta(n_1^* + i).$$

Let q is the smallest prime divisor of

$$(3.9) \quad c(n_1^*) - n_1^* + \delta(n_2) - 1 = (by (3.2)) \quad 2n_1^* + u + \delta(n_2) - 1.$$

Note that, since u is even, then q is odd. Let us prove that

$$(3.10) \quad k \geq \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k)).$$

In view of (3.7), for $i \leq k$, the number $h(n_1^* + i)$ divides $2n_1^* + u + \delta(n_1^* + i) - 1$.

Therefore, for $i = k$, we see that

$$(3.11) \quad h(n_1^* + k) \geq q.$$

Since, by (3.4), $h(n_1^* + k)$ divides $2k - u + 1 - 3\delta(n_1^* + k)$, then, using (3.11), we find

$$q \leq h(n_1^* + k) \leq 2k - u + 1 - 3\delta(n_1^* + k)$$

and (3.10) follows.

Now show that also

$$(3.12) \quad k \leq \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k)).$$

By the definition of k , for $1 \leq i < k$, we have $h(n_1^* + i) = 1$, and, using (3.10), we conclude that at least for $1 \leq i < \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + i))$ we have $h(n_1^* + i) = 1$. Show that $i = \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k))$ produces a nontrivial gcd. Indeed, according to (3.5), we have

$$\begin{aligned} & h(n_1^* + \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k))) = \\ & \gcd(n_1^* - \delta(n_1^* + k) + \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k)), \\ & 3n_1^* + u - 1 + \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k))) = \\ & \gcd(\frac{1}{2}((2n_1^* + u + \delta(n_1^* + k) - 1) + q), \\ (3.13) \quad & \frac{1}{2}(3(2n_1^* + u + \delta(n_1^* + k) - 1) + q)). \end{aligned}$$

From (3.10) and (3.13) it follows that q divides both of arguments of gcd.

Therefore,

$$h(n_1^* + \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k))) \geq q \geq 3.$$

Thus, by the definition of number n_2 , we have

$$k = n_2 - n_1^* \leq \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k)).$$

Therefore,

$$(3.14) \quad k = \frac{1}{2}(q + u - 1 + 3\delta(n_1^* + k)).$$

On the other hand, according to (3.8), $h(n_1^* + k)$, divides $2k - u + 1 - 3\delta(n_1^* + k)$, or, taking into account (3.14), divides q . Therefore,

$$(3.15) \quad h(n_2) = h(n_1^* + k) = q.$$

According to (3.14)-(3.15), we have

$$(3.16) \quad h(n_2) = q = 2k - u - 3\delta(n_2) + 1.$$

Nevertheless, by (2.8), $n_2 = 2n_1 + T_h - h - 3 \equiv 1 \pmod{2}$ and, by (3.3) $u = T_h - h - 2l_h - 6$. Therefore,

$$(3.17) \quad q = 2k - u - 5 = 2(n_2 - n_1^*) - u - 5 = 2n_1 + T_h - h - 5.$$

Thus $2n_1 + T_h - h - 5$ is prime. This completes proof of Theorem 1. ■

Corollary 2. *If $p_1 < p_2$ are consecutive greater of twin primes giving by Theorem 1, then $p_2 \geq 2p_1 - 1$.*

Proof. Since (see already proved observation 2) $n_2 - 3 \geq 2(n_1 - 3)$, then $q = n_2 - 2 \geq 2(n_1 - 2) - 1$, where, by the inductive supposition, $n_1 - 2$ is greater of twin primes. Now the corollary follows in the frameworks of the induction. ■

Corollary 3.

$$T_h \equiv h \pmod{6}.$$

Proof. The corollary immediately follows from the well known fact that the half-sum of twin primes not less than 5 is a multiple of 6. Therefore, $2n_1 + T_h - h - 6 \equiv 0 \pmod{6}$. Since, by the condition, $2n_1 \equiv 0 \pmod{6}$, then we obtain the corollary. ■

Now the observation 5) follows in the frameworks of the induction. The same we can say about observation 4).

The observed weak excesses of the exact estimate of Corollary 2 indicate to the smallness of T_h and confirm, by Theorem 1, Conjecture 1.

4. ESTIMATES OF RATIOS $c(n)/n$ AND STRONGER CONJECTURE

From the construction of Section 2 it easily follows that only in the fundamental points of the considered sequence we have $\gamma(n) := c(n)/n = 2$. Moreover, only in two points following after every fundamental point we have the values of $\gamma(n)$ less than 2. Namely, if n is a fundamental point, then in the point $\nu = n + 1$ we have $\gamma(\nu) = 2 - \frac{1}{\nu}$ and in the point $\mu = n + 2$ we have $\gamma(\mu) = 2 - \frac{2}{\mu}$. On the other hand, using induction, it is easy to

prove that

$$\frac{c(n)}{n} \leq \begin{cases} 3, & \text{if } n \text{ is even,} \\ 3 - \frac{6}{n}, & \text{if } n \text{ is odd.} \end{cases}$$

Indeed, let

$$c(n-1) \leq \begin{cases} 3(n-1), & \text{if } n \text{ is even,} \\ 3(n-1) - 6, & \text{if } n \text{ is odd.} \end{cases}$$

Since

$$h(n) = c(n) - c(n-1) \begin{cases} n \text{ and } c(n-1), & \text{if } n \text{ is even,} \\ n-2 \text{ and } c(n-1), & \text{if } n \text{ is odd,} \end{cases}$$

then

$$h(n) \leq \begin{cases} 3n - c(n-1), & \text{if } n \text{ is even,} \\ 3n - 6 - c(n-1), & \text{if } n \text{ is odd,} \end{cases}$$

and

$$c(n) = c(n-1) + h(n) \leq \begin{cases} 3(n-1), & \text{if } n \text{ is even,} \\ 3(n-1) - 6, & \text{if } n \text{ is odd.} \end{cases}$$

Thus we proved the following estimates.

Proposition 1.

$$(4.1) \quad 2 - \frac{2}{n-1} \leq \gamma(n) \leq \begin{cases} 3, & \text{if } n \text{ is even,} \\ 3 - \frac{6}{n}, & \text{if } n \text{ is odd.} \end{cases}$$

In points n of the main increments we have $\gamma(n) = 3 - \frac{6}{n}$. The first terms of the sequence $\{\beta_j\}$ for which $\gamma(\beta_j) = 3$ are:

$$18, 20, 66, 150, 156, 1326, 10904, 10908, 10910, \dots$$

It is easy to see that observation 6) one can replace by, e.g., the observation that, for every $i \geq 3$, in the maximal point $\rho^{(i)}$ of a nontrivial increment before fundamental point m_i we have

$$(4.2) \quad \gamma(\rho^{(i)}) \geq 2.5 \quad .$$

Indeed, putting in (2.5) $n_1 := n_{i-1} = m_{i-1} + 3$, $n_2 := n_i = m_i + 3$ and $n_1 + l_h := \rho^{(i)}$, such that, by (2.7) (see the second fundamental point of the inductive process), $T_h - h - 6 := m_i - 2n_{i-1}$ we, by the supposition, have

$$(4.3) \quad \gamma(\rho^{(i)}) = \frac{\rho^{(i)} + m_i}{\rho^{(i)}} \geq 2.5 \quad .$$

Thus

$$(4.4) \quad \rho^{(i)} \leq \frac{2}{3}m_i.$$

This means that the distance between $\rho^{(i)}$ and m_i is not less than $m_i/3$. Since we have $x/3 > \sqrt{x-1} + 4$, for $x \geq 30$, then observation 6) follows for $m_{i+1} \geq 30$.

Our stronger conjecture is the following.

Conjecture 3. *Let $m_{i-1} < m_i$ be adjacent fundamental points. Let $\rho^{(i)}$ be the maximal point of a nontrivial increment before m_i . Then*

$$(4.5) \quad \lim_{i \rightarrow \infty} \frac{\rho^{(i)}}{m_i} = \frac{1}{2}.$$

5. A SUFFICIENT CONDITION FOR THE INFINITY OF TWIN PRIMES

Put

$$(5.1) \quad \rho^{(i)} = \lambda_i n_{i-1}, \quad \lambda_i \geq 1.$$

Conjecture 4. *For every $i \geq 6$, $\lambda_i \leq 5/4$.*

Theorem 4. *If Conjecture 4 is true, then we have infinitely many twin primes.*

Proof. Since

$$(5.2) \quad c(\lambda_i n_{i-1}) = \gamma(\rho^{(i)}) \lambda_i n_{i-1},$$

then the distance r_i between $\rho^{(i)}$ and m_i is defined by the equation

$$(5.3) \quad \frac{\gamma(\rho^{(i)}) \lambda_i n_{i-1} + r_i}{\lambda_i n_{i-1} + r_i} = 2.$$

Thus

$$(5.4) \quad r_i = \gamma(\rho^{(i)}) \lambda_i n_{i-1} - 2 \lambda_i n_{i-1}$$

and we have

$$(5.5) \quad m_i = \lambda_i n_{i-1} + r_i = \lambda_i n_{i-1} (\gamma(\rho^{(i)}) - 1).$$

Put

$$(5.6) \quad m_i = \lambda_i n_{i-1} (\gamma(\rho^{(i)}) - 1) = (2 + \mu_i) n_{i-1}.$$

Since, by Theorem 2, which was proved independently from observation 6), we have

$$2 \leq \frac{m_i}{m_{i-1}} = \frac{m_i}{n_{i-1} - 3},$$

then $m_i \geq 2n_{i-1} - 6$ and, by (5.6),

$$(5.7) \quad 2 + \mu_i = \lambda_i (\gamma(\rho^{(i)}) - 1) \geq 2 - \frac{6}{n_{i-1}}.$$

Furthermore, by the condition, $\lambda_i \leq 5/4$. Therefore, we have

$$5/4 \geq \lambda_i \geq (2 - \frac{6}{n_{i-1}}) \frac{1}{\gamma(\rho^{(i)}) - 1}.$$

Note that, for $i \geq 6$, we have $n_{i-1} \geq 141$. Therefore,

$$(5.8) \quad \gamma(\rho^{(i)}) \geq 1 + \frac{4}{5}(2 - \frac{6}{n_{i-1}}) \geq 5/2.$$

By (4.3)-(4.4), this means that observation 6) follows and the numbers $m_i \mp 1$ are twin primes.

On the other hand, by (5.5) and Proposition 1, we have

$$(5.9) \quad m_i = \lambda n_{i-1} (\gamma(\rho^{(i)}) - 1) \leq 2\lambda_i n_{i-1} \leq 2.5n_{i-1}.$$

The latter inequality ensures the infinity of the fundamental points of the considered sequence and, consequently, the infinity of twin primes. ■

Moreover, if Conjecture 4 is true, then verifying a finite set of integers beginning with $n = 2$, from Theorem 4 we obtain that:

Between $n \geq 2$ and $3n$ we have at least one pair of twin primes.

Note that, the first real values of $\lambda_i = \frac{\rho^{(i)}}{n_{i-1}}$, $i \geq 6$ are:

$$\begin{array}{lll} \frac{156}{141} = 1.106...; & \frac{348}{315} = 1.104...; & \frac{661}{661} = 1.000...; \\ \frac{1339}{1323} = 1.012...; & \frac{2712}{2661} = 1.019...; & \frac{5496}{5421} = 1.013... , \dots \end{array}$$

Note that if the last miner increment $\rho^{(i)}$ after the point of the main increment n_{i-1} is known, then the following miner increment is

$$(5.10) \quad n_i = c(\rho^{(i)}) - \rho^{(i)}.$$

It easy follows from (2.5)-(2.6).

6. TO EVERY INTEGER $m \geq 4$ CORRESPONDS A PAIR OF TWIN PRIMES

$(p, p+2)$ SUCH THAT $p+2 \geq m$

Given $m \geq 4$, we give a very simple rule to calculate a pair of twin primes $(p, p+2)$ such that $p+2 \geq m$. Although till now we are able to prove a private case of this rule, we absolutely do not doubt that it is always true! For every positive integer m , consider the following sequence:

$$c^{(m)}(1) = m; \text{ for } n \geq 2,$$

$$(6.1) \quad c^{(m)}(n) = c^{(m)}(n-1) + \begin{cases} \gcd(n, c^{(m)}(n-1)), & \text{if } n \text{ is even} \\ \gcd(n-2, c^{(m)}(n-1)), & \text{if } n \text{ is odd.} \end{cases}$$

Thus for every m this sequence has the the same formula that the considered one but another initial condition. Our astonishing observation is the following.

Conjecture 5. *Let n^* , where $n^* = n^*(m)$, be point of the last nontrivial increment of $\{c^{(m)}(n)\}$ on the set $N_m = \{1, \dots, m\}$ and $n^* = 1$, if there is not any nontrivial increment on N_m . Then numbers $c^{(m)}(n^*) - n^* \mp 1$ are twin primes.*

Evidently, $c^{(m)}(n^*) - n^* + 1 \geq m$ and the equality holds if and only if $n^* = 1$.

Example 2. *Let $m = 20$. Then $n^* = 12$ and $c^{(m)}(n^*) = 42$. Thus numbers $42 - 12 \mp 1$ are twin primes (29, 31).*

Example 3. *Let $m = 577$. Then $n^* = 156$ and $c^{(m)}(n^*) = 1038$. Thus numbers $1038 - 156 \mp 1$ are twin primes (881, 883).*

Example 4. *Let $m = 3000$. Then $n^* = 2$ and $c^{(m)}(n^*) = 3002$. Thus numbers $3002 - 2 \mp 1$ are twin primes (2999, 3001).*

The case of $n^* = 1$ we formulate as the following criterion.

Criterion 1. *A positive integer $m > 3$ is a greater of twin primes if and only if the points $1, \dots, m$ are points of trivial increments of sequence $\{c^{(m)}(n)\}$.*

Proof. By the condition,

$$c^{(m)}(1+i) = m+i, \quad c^{(m)}(2+i) = m+i+1,$$

Therefore, if i is even, then

$$\gcd(2+i, m+i) = 1,$$

or

$$\gcd(m-2, i+2) = 1.$$

If i is odd, then

$$\gcd(i, m+i) = 1,$$

or

$$\gcd(m, i) = 1.$$

Since i is arbitrary from N_m , then both of numbers $m-2$, m are primes. The converse statement is also evident. ■

7. A THEOREM ON TWIN PRIMES WHICH IS INDEPENDENT ON OBSERVATION OF TYPE 6)

Here we present a new sequence $\{a(n)\}$ with the quite analogous definition of fundamental and miner points for which Corollary 1 is true in a stronger formulation. Using a construction close to those ones that we considered in [3], consider the sequence defined as the following:

$$(7.1) \quad a(180) = 360 \text{ and for } n \geq 181, \\ a(n) = \begin{cases} a(n-1) + 1, & \text{if } \gcd(n + (-1)^n - 1, a(n-1)) = 1; \\ 3(n-2) & \text{otherwise} \end{cases}.$$

Definition 1. A point m_i is called a fundamental point of sequence (7.1), if it has the form $m_i = 6t$ and $a(m_i) = 2m_i$. The increments in the points $m_i + 3$ we call the main increments. Other nontrivial increments we call miner increments.

The first fundamental point of sequence (7.1) is $m_1 = 180$.

Theorem 5. If the sequence $\{a(n)\}$ contains infinitely many fundamental points, then there exist infinitely many twin primes.

Proof. We use induction. Note that numbers $m_1 \mp 1$ are twin primes: 179 and 181. Suppose that, for some $i \geq 1$, the numbers $m_i \mp 1$ are twin primes. Put $n_i = m_i + 3$. Then $n_i \equiv 3 \pmod{6}$ and we have

$$\begin{aligned} a(n_i - 3) &= 2n_i - 6 \\ a(n_i - 2) &= 2n_i - 5, \\ a(n_i - 1) &= 2n_i - 4, \\ a(n_i) &= 3n_i - 6, \end{aligned}$$

We see that the main increment is $n_i - 2$. By the condition, before m_{i+1} we can have only a finite set of miner increments. Suppose that, they are in the points $n_i + l_j, j = 1, \dots, h_i$. Then, by (7.1), we have

$$a(n_i + 1) = 3n_i - 5,$$

...

$$\begin{aligned}
a(n_i + l_1 - 1) &= 3n_i + l_1 - 7, \\
a(n_i + l_1) &= 3n_i + 3l_1 - 6, \\
&\dots \\
a(n_i + l_2 - 1) &= 3n_i + 2l_1 + l_2 - 7, \\
a(n_i + l_2) &= 3n_i + 3l_2 - 6, \\
&\dots \\
a(n_i + l_h - 1) &= 3n_i + 2l_{h-1} + l_h - 7, \\
(7.2) \quad a(n_i + l_h) &= 3n_i + 3l_h - 6, \\
&\dots
\end{aligned}$$

$$\begin{aligned}
(7.3) \quad a(n_{i+1} - 3) &= 2n_{i+1} - 6 \\
a(n_{i+1} - 2) &= 2n_{i+1} - 5, \\
a(n_{i+1} - 1) &= 2n_{i+1} - 4,
\end{aligned}$$

$$(7.4) \quad a(n_{i+1}) = 3n_{i+1} - 6.$$

Note that, in every step from (7.2) up to (7.3) we add 1 simultaneously to values of the arguments and of the right hand sides. Thus in the fundamental point $m_{i+1} = n_{i+1} - 3$ we have

$$n_i + l_h + x = n_{i+1} - 3$$

and

$$3n_i + 3l_h - 6 + x = 2n_{i+1} - 6$$

such that

$$n_{i+1} = 2n_i + 2l_h - 3.$$

Now we should prove that the numbers

$$n_{i+1} - 4 = 2n_i + 2l_h - 7, \quad n_{i+1} - 2 = 2n_i + 2l_h - 5$$

are twin primes. We have

$$\begin{aligned}
a(n_i + l_h + t) &= 3n_i + 3l_h - 6 + t, \\
(7.5) \quad a(n_i + l_h + t + 1) &= 3n_i + 3l_h - 5 + t,
\end{aligned}$$

where $0 \leq t \leq n_i + l_h - 6$. Distinguish two case.

1) Let l_h be even. Then, for even values of t the numbers $n_i + l_h + t + 1$ are even and from equalities (7.5) we have

$$\gcd(n_i + l_h + t + 1, 3n_i + 3l_h - 6 + t) = 1.$$

It is easy to see that $l_h + 1$ is not multiple of 3. Indeed, it is sufficient to choose $t = 6$. Thus $2l_h - 1$ is not multiple of 3 and, therefore, $N = 2n_i + 2l_h - 7$ also is not multiple of 3.

Furthermore, considering t not multiple of 3, from equalities (7.5) we have

$$\gcd(3n_i + 3l_h + 3t + 3, 3n_i + 3l_h - 6 + t) = 1$$

and

$$\gcd(2t + 9, 2n_i + 2l_h - 7) = 1, \quad 0 \leq t \leq n_i + l_h - 6, \quad t \equiv 2, 4 \pmod{6}.$$

Now in order to prove that N is prime it is sufficient to use t of the form $t = 6u + 2$. Since $0 \leq t \leq n_i + l_h - 6$, then $9 \leq 2t + 9 = 12u + 13 \leq 2n_i + 2l_h - 3$ and $0 \leq u \leq (n_i + l_h - 8)/6 = (N - 9)/12 > (n_i - 8)/6$. Note that, for the considered values of $n_i (\geq 183)$ we have $\frac{n_i - 8}{6} > \sqrt{2n_i - 16}$. Therefore, $(N - 9)/12 > \sqrt{N}$. Let $p \leq \sqrt{N}$. Consider the congruence $12u + 13 \equiv 0 \pmod{p}$. Choose a solution $u \in \{0, 1, \dots, p-1\}$. Then $u \leq \sqrt{N} < (N - 9)/12$ and we conclude that $\gcd(N, p) = 1$. Thus N is prime.

On the other hand, for odd values of t , taking into account that numbers $n_i + l_h + t + 1$ are odd, from equalities (7.5) we have

$$\gcd(n_i + l_h + t - 1, 3n_i + 3l_h - 6 + t) = 1.$$

Note that $l_h - 1$ is not multiple of 3. Indeed, it is sufficient to choose $t = 3$. Thus $2l_h - 5$ is not multiple of 3 and, therefore, $M = 2n_i + 2l_h - 5$ also is not multiple of 3.

Let now t is not multiple of 3. Then

$$\gcd(3n_i + 3l_h + 3t - 3, 3n_i + 3l_h - 6 + t) = 1$$

and

$$\gcd(2t + 3, 2n_i + 2l_h - 5) = 1, \quad 0 \leq t \leq n_i + l_h - 6, \quad t \equiv \pm 1 \pmod{6}.$$

In order to prove that M is prime it is sufficient to use t of the form $t = 6u + 1$. Since $0 \leq t \leq n_i + l_h - 6$, then $3 \leq 2t + 3 = 12u + 5 \leq 2n_i + 2l_h - 9$ and $0 \leq u \leq (n_i + l_h - 7)/6 = (M - 9)/12 > (n_i - 7)/6$ and exactly as for N we obtain that M is prime as well and the numbers N and M are twin primes.

2) Let l_h be odd. Then, using again equalities (7.5), by the same way, we show that the numbers N, M are twin primes. This completes the induction. ■

Acknowledgment. The author is grateful to Daniel Berend (Ben Gurion University, Israel) for useful discussions; he also is grateful to Richard Mathar (Leiden University, Netherlands) and Konstantin Shukhmin (Dunedin, New Zealand) for an important help in the numerical calculations.

REFERENCES

- [1] . E. S. Rowland *A natural prime-generating recurrence* J.Integer Seq., v.11(2008), Article 08.2.8
- [2] . V. Shevelev, *An infinite set of generators of primes based on the Roland idea and conjectures concerning twin primes*, <http://www.arxiv.org/abs/0910.4676> [math. NT].
- [3] . V. Shevelev, *Generalizations of the Rowland theorem*, <http://www.arxiv.org/abs/0911.3491> [math. NT].
- [4] . N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences* ([http : //www.research.att.com/~njas](http://www.research.att.com/~njas))

DEPARTMENTS OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA 84105, ISRAEL. E-MAIL:SHEVELEV@BGU.AC.IL